



Organization for angioedema
Bramerveld 53, 2151 LA, Nieuw-Vennep
+316 10517525
info@hae-qe.nl
KvK: 40413239

Mrs

has hereditary angioedema (congenital C1-inhibitor deficiency).

Treatment with C1-inhibitor concentrate (Cynrize[®], 1000 U iv) or icatibant (Firazyr[®], 30 mg subcutaneously) is urgently warranted in case of an attack of angioedema. She may carry this treatment (including equipment for the intravenous administration of this drug (needles, syringes, etc.)). She may need some help with the administration.

If the concentrate is not readily available, administration of any other iv C1-inhibitor concentrate (Conestat alfa/Ruconest[®] or Berinert[®]) or icatibant (Firazyr[®]) 30 mg subcutaneously may serve as an alternative. Should none of these products be available, then plasma (large quantities, i.e. 6 units i.v.) may effectively reduce the attack burden. Steroids and epinephrine are not effective.



Organization for angioedema
Bramerveld 53, 2151 LA, Nieuw-Vennep
+316 10517525
info@hae-qe.nl
KvK: 40413239

Mr

has hereditary angioedema (congenital C1-inhibitor deficiency).

Treatment with C1-inhibitor concentrate (Cynrize[®], 1000 U iv) or icatibant (Firazyr[®], 30 mg subcutaneously) is urgently warranted in case of an attack of angioedema. He may carry this treatment (including equipment for the intravenous administration of this drug (needles, syringes, etc.)). He may need some help with the administration.

If the concentrate is not readily available, administration of any other iv C1-inhibitor concentrate (Conestat alfa/Ruconest[®] or Berinert[®]) or icatibant (Firazyr[®]) 30 mg subcutaneously may serve as an alternative. Should none of these products be available, then plasma (large quantities, i.e. 6 units i.v.) may effectively reduce the attack burden. Steroids and epinephrine are not effective.



Organization for angioedema
Bramerveld 53, 2151 LA, Nieuw-Vennep
+316 10517525
info@hae-qe.nl
KvK: 40413239

Child

has hereditary angioedema (congenital C1-inhibitor deficiency).

Treatment with C1-inhibitor concentrate (Cynrize[®], 1000 U iv) or icatibant (Firazyr[®], 30 mg subcutaneously) is urgently warranted in case of an attack of angioedema. {He/she} may carry this treatment (including equipment for the intravenous administration of this drug (needles, syringes, etc.)). {He/she} may need some help with the administration.

If the concentrate is not readily available, administration of any other iv C1-inhibitor concentrate (Conestat alfa/Ruconest[®] or Berinert[®]) or icatibant (Firazyr[®]) 30 mg subcutaneously may serve as an alternative. Should none of these products be available, then plasma (large quantities, i.e. 6 units i.v.) may effectively reduce the attack burden. Steroids and epinephrine are not effective.